

EQUIVARIANT UNFOLDINGS OF G -STRATIFIED PSEUDOMANIFOLDS

DALMAGRO F.

To Rodolfo Ricabarra, in memoriam.

ABSTRACT. For any abelian compact Lie group G , we introduce a family of G -stratified pseudomanifolds, whose main feature is the preservation of the orbit spaces in the category of stratified pseudomanifolds. Which generalize a previous definition given in [8]. We also find a sufficient condition for the existence of equivariant unfoldings, so we have Intersection Cohomology with differential forms, as defined in [9]. Moreover, if G act on a manifold M , we find a equivariant unfolding of M which induce a canonical unfolding on the k -orbits space for every closed subgroup K of G .

INTRODUCTION

Let X be a Thom-Mather stratified space with depth $d(X) = n$. The De Rham Intersection Cohomology of X with differential forms was defined in [3] by means of an auxiliar construction called *unfolding*, which is a continuous map $L : \tilde{X} \rightarrow X$ where \tilde{X} is a smooth manifold obtained trough a finite composition

$$L : \tilde{X} = X_n \xrightarrow{L_n} X_{n-1} \rightarrow \cdots \rightarrow X_1 \xrightarrow{L_1} X_0 = X$$

of topological operations $X_i \xrightarrow{L_i} X_{i-1}$, called *elementary unfoldings*. This iterative construction is possible because the stratification of X is controlled by the existence of a family of conical fiber bundles over the singular strata. Later in [9] we find a more abstract definition of unfoldings, which impose some conditions of transversality over the singular strata. For instance, if the depth of X is 1 then the first elementary unfolding of X is an unfolding in the new sense.

Now let G be a compact Lie group. We introduce the definition of a G -stratified pseudomanifold in the category of stratified pseudomanifolds. Our definition is related to a previous one given in [8]. A G -stratified pseudomanifold is a stratified pseudomanifold in the usual sense together with a continuous action preserving the strata, and whose local model near each singular strata is given by a conical slice. We also give a definition of equivariant unfoldings, which is a suitable adaptation

Date: July 2003.

1991 Mathematics Subject Classification. 35S35; 55N33.

Key words and phrases. Intersection Cohomology, Stratified Pseudomanifolds.

of the usual definition of unfolding to the family of G -stratified pseudomanifolds. We give a sufficient condition for the existence of equivariant unfoldings, which is related to the choice of a good family of tubular neighborhoods and a sequence of equivariant elementary unfoldings of X . Each elementary unfolding induces an elementary unfolding on each orbit space together with a factorization diagram.

The content of this paper is as follows:

In §1 we introduce the category of stratified pseudomanifolds.

In §2 we define stratified G -pseudomanifolds and study their corresponding K -orbit spaces, for a closed subgroup K of G .

In §3 we study equivariant tubular neighborhoods, which are equivariant versions of the usual ones.

In §4 we introduce the concept of an equivariant explosion of a G -pseudomanifold, and present elementary explosions as given in [1].

In §5 we define G -pseudomanifolds X stratified with a transversal Thom-Mather structure whose main feature is that the sequence of elementary explosions determine an equivariant explosion of the space X , which project naturally onto an equivariant explosion of X/K , for a closed subgroup K of G .

1. STRATIFIED PSEUDOMANIFOLDS

In this section we review the usual definitions of stratified spaces, stratified morphisms and stratified pseudomanifolds. For a more detailed introduction see [5], [7].

1.1. Stratified spaces Let X be a Hausdorff, locally compact and 2nd countable space. A **stratification** of X is a locally finite partition \mathcal{S}_X satisfying:

- (i) Each element $S \in \mathcal{S}_X$ is a connected manifold with the induced topology, which is a **stratum** of X .
- (ii) If $S' \cap \overline{S} \neq \emptyset$ then $S' \subset \overline{S}$ for any two strata $S, S' \in \mathcal{S}_X$. In this case we write $S' \leq S$ and we say that S **incides** on S' .

We say that (X, \mathcal{S}_X) is a **stratified space** whenever \mathcal{S}_X is a stratification of X .

With the above conditions, the incidence relationship is a partial order on \mathcal{S}_X . More over, since \mathcal{S}_X is locally finite, any strictly ordered chain

$$S_0 < S_1 < \cdots < S_m$$

in \mathcal{S}_X is finite. The **depth** of X is by definition the supremum (possibly infinite) of the integers m such that there is a strictly ordered chain as above. We write this as $d(X)$.

The maximal (resp. minimal) strata in X are open (resp. closed) in X . A **singular** stratum is a non-maximal stratum in X . The union of the singular strata is the **singular part** of X , denoted by $\Sigma \subset X$, which is closed in X . Its complement $X - \Sigma$ is open and dense in X . The family of minimal strata will often be denoted

by \mathcal{S}_X^{\min} , while the union of minimal strata will be denoted by Σ^{\min} , which we call the **minimal part** of X .

1.2. Examples Here there are some examples of stratified spaces.

- (1) For any manifold M the trivial stratification of M is the family

$$\mathcal{S}_M = \{C : C \text{ is a connected component of } M\}$$

- (2) For any connected manifold M , the space $M \times X$ is a stratified space, with the stratification

$$\mathcal{S}_{M \times X} = \{M \times S : S \in \mathcal{S}_X\}$$

Notice that $d(M \times X) = d(X)$.

- (3) The **cone** of a compact stratified space L is the quotient space

$$c(L) = L \times [0, \infty) / L \times \{0\}$$

We write $[p, r]$ for the equivalence class of $(p, r) \in L \times [0, \infty)$. The symbol $*$ will be used for the equivalence class of $L \times \{0\}$, this is the **vertex** of the cone. The family

$$\mathcal{S}_{c(L)} = \{*\} \cup \{S \times (0, \infty) : S \in \mathcal{S}_L\}$$

is the canonical stratification of $c(L)$. Notice that $d(c(L)) = d(L) + 1$.

1.3. Stratified subspaces and morphisms Let (X, \mathcal{S}_X) be a stratified space. For each subset $Z \subset X$ the **induced partition** is the family

$$\mathcal{S}_{Z/Y} = \{C : C \text{ is a connected component of } Z \cap S, S \in \mathcal{S}_X\}$$

We will say that Z is a **stratified subspace** of X , whenever the induced partition on Z is a stratification of Z .

Now let (Y, \mathcal{S}_Y) be another stratified space. A **morphism** (resp. **isomorphism**) is a continuous map $f : X \rightarrow Y$ (resp. homeomorphism) which smoothly (resp. diffeomorphically) sends strata into strata. In particular, f is a **embedding** if $f(X)$ is a stratified subspace of Y and $f : X \rightarrow f(X)$ is an isomorphism.

Henceforth, we will write $\text{Iso}(X, \mathcal{S}_X)$ for the group of isomorphisms of a stratified space X . The following statement will be used later, we leave the proof to the reader.

Lemma 1.4. *Let (X, \mathcal{S}_X) be a stratified space, and $\mathfrak{F} \subset \mathcal{S}_X$ a subfamily of equidimensional strata. The connected components of $M = \bigcup_{S \in \mathfrak{F}} S$ are the strata in \mathfrak{F} .*

Stratified pseudomanifolds were used by Goresky and MacPherson in order to introduce the Intersection Homology and extend the Poincaré duality to the family of stratified spaces. For a brief introduction the reader can see [5].

1.5. Stratified pseudomanifolds The definition of a stratified pseudomanifold is made by induction on the depth of the space. More precisely:

- (1) A stratified pseudomanifold of depth 0 is a manifold with the trivial stratification.
- (2) An arbitrary stratified space (X, \mathcal{S}_X) is a **stratified pseudomanifold** if, for any singular stratum $S \in \mathcal{S}_X$, there is a compact stratified pseudomanifold L_S depending on S (called a **link** of S) such that each point $x \in S$ has a coordinate neighborhood $U \subset S$ and an embedding onto an open subset of X .

$$\varphi : U \times c(L_S) \rightarrow X$$

such that $x \in \text{Im}(\varphi)$. The pair (U, φ) is a **chart** of x **modelled** on L_S .

1.6. Examples Here are some examples of stratified pseudomanifolds.

- (1) If X is a stratified pseudomanifold, then any open subset $A \subset X$ is also a stratified pseudomanifold. Also the product $M \times X$ (with the canonical stratification) is a stratified pseudomanifold, for any manifold M .
- (2) If L is a compact stratified pseudomanifold, then $c(L)$ is a stratified pseudomanifold.

2. G -STRATIFIED PSEUDOMANIFOLDS

From now on, we fix an abelian compact Lie group G . We will study the family of actions of G which preserve the strata. Our definition is strongly related to the previous one given in [8]. Also some easy proofs in this section can be seen in [6].

Given a stratified space (X, \mathcal{S}_X) and a effective action $\Phi : G \times X \rightarrow X$; we write $\Phi(g, x) = gx$ for any $g \in G$, $x \in X$. We denoted X/K by the K -orbit space for every K closed subgroup of G , and by $\pi : X \rightarrow X/K$ the orbit map. The group of G -equivariant isomorphisms of X will be denoted by $\text{Iso}_G(X, \mathcal{S}_X)$.

2.1. G -stratified spaces We say that X is **G -stratified** whenever:

- (1) For each stratum $S \in \mathcal{S}_X$ the points of S all have the same isotropy group, denoted by G_S .
- (2) Each $g \in G$ induces an isomorphism $\Phi_g : X \rightarrow X \in \text{Iso}_G(X, \mathcal{S}_X)$.

The orbit space X/K inherits a **canonical stratification** given by the family

$$\mathcal{S}_{X/K} = \{\pi(S) : S \in \mathcal{S}_X\}$$

Notice also that $d(X) = d(X/K)$.

2.2. Examples Here are some examples of G -stratified spaces:

- (1) Each G -manifold M has a natural structure of G -stratified space, when M is endowed with the stratification given by orbit types.
- (2) If X is a G -stratified space, then $M \times X$ is a G -stratified space with the action $g(m, x) = (m, gx)$; for any manifold M .
- (3) If L is a compact G -stratified space then $c(L)$, with the action $g[x, r] = [gx, r]$, is a G -stratified space.

Lemma 2.3. *Let G be an abelian compact Lie group, $K \subset G$ a closed subgroup. Then for any G -stratified space X the orbit space X/K is a G/K -stratified space.*

Proof. Write $\bar{g} \in G/K$ for the equivalence class of $g \in G$. Consider the quotient action

$$\bar{\Phi} : G/K \times X/K \rightarrow X/K \quad \bar{g} \cdot \pi(x) = \pi(gx)$$

This action is well defined because G is abelian. So:

• **The isotropy groups are constant over the strata of X/K :** This is straightforward, since for each stratum $S \in \mathcal{S}_X$ we have

$$(G/K)_{\pi(S)} = KG_S/K$$

Hence $\pi(S)$ has constant isotropy.

• **Each \bar{g} induces an isomorphism $\bar{\Phi}_g \in G/K \in \text{Iso}(X/K, \mathcal{S}_{X/K})$:** For each $g \in G$ we have a K -equivariant isomorphism $\Phi_g \in \text{Iso}(X, \mathcal{S}_X)$. Passing to the quotients we obtain an isomorphism $\bar{\Phi}_g \in G/K \in \text{Iso}(X/K, \mathcal{S}_{X/K})$. The differentiability of this map on $\pi(S)$ is immediate from the following commutative diagram

$$\begin{array}{ccc} S & \xrightarrow{\Phi_g} & gS \\ \pi \downarrow & & \downarrow \pi \\ \pi(S) & \xrightarrow{\bar{\Phi}_g} & \pi(gS) \end{array}$$

□

Now we introduce the definition of a G -stratified pseudomanifold.

2.4. G -stratified pseudomanifolds A G -stratified pseudomanifold is a stratified pseudomanifold in the usual sense, endowed with a structure of G -stratified space (i.e. G acts by isomorphisms) and whose local model is described through conical slices. Conical slices were introduced in [8] in order to state a sufficient condition on any continuous action of a compact Lie group (abelian or not) on stratified pseudomanifold so that the corresponding orbit space would remain in the same class of spaces.

Let (X, \mathcal{S}_X) be a G -stratified space. Take a singular stratum $S \in \mathcal{S}_X$ a point $x \in S$. A **conical slice** of x in X is a slice S_x in the usual sense of [2], with a conical part transverse to the stratum S . In other words:

- (1) S_x is an invariant G_S -space containing x .
- (2) For any $g \in G$, if $gS_x \cap S_x \neq \emptyset$ then $g \in G_S$.
- (3) $G_S S_x$ is open in X . And
- (4) There is a G_S -equivalence $\beta : \mathbb{R}^i \times c(L) \rightarrow S_x$ where $i \geq 0$ and L is a compact G_S -stratified space. Here the action of G_S on \mathbb{R}^i is trivial (notice that β induces on S_x a structure of G_S -stratified space).

The definition of a **G -stratified pseudomanifold** is made by induction on the depth of the space. A G -stratified pseudomanifold with depth 0 is a manifold with a smooth free action of G . In general, we will say that X is a G -stratified pseudomanifold if, for each singular stratum $S \in \mathcal{S}_X$, there is a compact G_S -stratified pseudomanifold L_S such that each point $x \in S$ has a conical slice

$$\beta : \mathbb{R}^i \times c(L_S) \rightarrow S_x$$

and the usual map on the twisted product

$$\alpha : G \times_{G_S} S_x \rightarrow X \quad \alpha([g, y]) = gy$$

is an equivariant (stratified) embedding on an open subset of X . We say that the triple (S_x, β, L_S) is a **distinguished slice** of x .

2.5. Examples Here there are some examples of G -stratified pseudomanifolds.

- (1) Take a smooth effective action $\Phi : G \times M \rightarrow M$ with fixed points on a manifold M endowed with the stratification by orbit types. By the Equivariant Slice Theorem, M is a G -stratified pseudomanifold.
- (2) If X is a G -stratified pseudomanifold then $M \times X$ is a G -stratified pseudomanifold with the obvious action.
- (3) If L is a compact G -stratified pseudomanifold, then $c(L)$ is a G -stratified pseudomanifold with the obvious action.
- (4) Any invariant open subspace of a G -stratified pseudomanifold is itself a G -stratified pseudomanifold.

♣ *Remark 2.6. Each G -stratified pseudomanifold is a stratified pseudomanifold in the previous sense.*

To see this, proceed by induction on the depth. Take a G -stratified pseudomanifold X . For $d(X) = 0$ the statement is trivial. Assume the inductive hypothesis and suppose that $d(X) > 0$. Take a singular stratum $S \in \mathcal{S}_X$, a point $x \in S$ and a distinguished slice (S_x, β, L_S) of x . The isotropy subgroup G_S acts on G by the restriction of the group operation. We fix a slice S_e of the identity element $e \in G$ with respect to this action. Since $G_S S_e$ is open in G , the composition

$$\begin{aligned} (S_e \times \mathbb{R}^i) \times c(L_S) &\rightarrow S_e \times (\mathbb{R}^i \times c(L_S)) \rightarrow S_e \times S_x \rightarrow \\ &\rightarrow S_e \times (G_S \times_{G_S} S_x) \rightarrow (G_S S_e) \times_{G_S} S_x \rightarrow X \end{aligned}$$

is an embedding. Notice that L_S is a stratified pseudomanifold by induction. Since $S_e \times \mathbb{R}^i \simeq S_e G_S(S \cap S_x)$ is open in S . We have obtained a chart of x modelled on L_S .

♣ *Remark 2.7. If X is a G -stratified pseudomanifold and K is any closed subgroup of G , then X is also a K -stratified pseudomanifold.*

It is straightforward that X is a K -stratified space. For any singular stratum S and any $x \in S$, in order to choose a distinguished slice in x we proceed as follows: Take a distinguished slice $\beta : \mathbb{R}^i \times c(L_S) \rightarrow S_x$ in x with respect to the action of G . Take also a slice V_e of the identity element $e \in G$ with respect to the action of

$G_S K$ in G . Then $\iota \times \beta : (V \times \mathbb{R}^i) \times c(L_S) \rightarrow VS_x$ is a distinguished slice of x with respect to the action of K .

Now we study the factorization of a G -stratified pseudomanifold when considered as a K -stratified pseudomanifold for any closed subgroup $K \subset G$.

Proposition 2.8. *Let G be a compact, abelian Lie group; $K \subset G$ a closed subgroup. If X is a G -stratified pseudomanifold then X/K is a G/K -stratified pseudomanifold.*

Proof. As before, write $\pi : X \rightarrow X/K$ for the orbit map induced by the action of K on X . Proceed by induction on $l = d(X)$. For $l = 0$ it is straightforward, since $d(X/K) = d(X) = 0$. Assume the inductive hypothesis and suppose that $d(X) > 0$. By §2.3, X/K is a G/K -stratified space, so we must verify the existence of conical slices.

Take a singular stratum $S \in \mathcal{S}_X$, fix a point $x \in S$ and a distinguished slice (S_x, β, L_S) of a x . The K -equivariant isomorphism $\beta : \mathbb{R}^i \times c(L_S) \rightarrow S_x$ induces an isomorphism on the orbit spaces

$$\bar{\beta} : \mathbb{R}^i \times c(L_S/G_S \cap K) \rightarrow \pi(S_x) \quad \bar{\beta}(b, [\bar{l}, r]) = \pi(\beta(b, [l, r]))$$

Now we will show that the triple $(\pi(S_x), \bar{\beta}, L_S/G_S \cap K)$ is a distinguished slice of $\pi(x) \in X/K$. We do it in three steps.

- **$\pi(S_x)$ is a slice of $\pi(x)$:** This is straightforward, since $(G/K)_{\pi(x)} = KG_S/K$, the quotient $\pi(S_x)$ is a $(G/K)_{\pi(x)}$ -space with the quotient action and the orbit map π is an open map.
- **$\bar{\beta}$ is a KG_S/K -equivalence:** This is immediate, since β is an H -equivalence. Notice that, by induction on the depth, $L_S/G_S \cap K$ is a KG_S/K -stratified pseudomanifold.
- **The induced map $\bar{\alpha} : (G/K) \times_{(G_S/G_S \cap K)} \pi(S_x) \rightarrow X/K$ is an embedding:** This $\bar{\alpha}$ is given by the rule $\bar{\alpha}([\bar{g}, \pi(z)]) = \bar{g} \cdot \pi(z)$, and is a homeomorphism. We consider the following commutative diagram

$$\begin{array}{ccc} G \times_{G_S} S_x & \xrightarrow{\alpha} & X \\ \pi \downarrow & & \downarrow \pi \\ (G/K) \times_{(G_S/G_S \cap K)} \pi(S_x) & \xrightarrow{\bar{\alpha}} & X/K \end{array}$$

Since the vertical arrows are submersions, and α is an embedding, we obtain that $\bar{\alpha}$ is an embedding. \square

3. TUBULAR NEIGHBORHOODS

Henceforth we fix a compact, abelian Lie group G , and a G -stratified pseudomanifold X . In this section we will study the family of equivariant tubular neighborhoods, which are equivariant version of the usual ones. Given a singular stratum S in X , a tubular neighborhood is just a locally trivial fiber bundle over a S , whose

fiber is $c(L_S)$, the cone of the link of S ; and whose structure group is $\text{Iso}_{G_S}(L_S, \mathcal{S}_{L_S})$.

We will clarify these ideas immediately, considering a previous step in our way: the definition of a G -stratified fiber bundle. The reader will find in [10] a detailed introduction to the fiber bundles, while [11] provides the usual definition of a tubular neighborhood in the stratified context (see also [2] for the smooth case).

3.1. G -stratified fiber bundles Let $\xi = (E, p, B, F)$ be a locally trivial fiber bundle with (maximal) trivializing atlas \mathcal{A} . We will say that ξ is a **G -stratified** whenever:

- (1) The total space E is a G -stratified space.
- (2) The base space B is a manifold, endowed with a smooth action $\Psi : G \times B \rightarrow B$ and with constant isotropy $H \subset G$ at all its points.
- (3) The fiber F is a H -stratified space.
- (4) The projection $p : E \rightarrow B$ is G -equivariant.
- (5) The group G acts by isomorphisms. In other words, each chart

$$\varphi : U \times F \rightarrow p^{-1}(U) \in \mathcal{A}$$

is H -equivariant; and for any two charts $(U, \varphi), (U', \varphi') \in \mathcal{A}$ such that $U' \cap g^{-1}U \neq \emptyset$ for some $g \in G$, there is a map

$$g_{\varphi, \varphi'} : U' \cap g^{-1}U \rightarrow \text{Iso}_H(F, \mathcal{S}_F)$$

such that

$$\varphi^{-1}g\varphi'(b, z) = (gb, g_{\varphi, \varphi'}(b)z)$$

Lemma 3.2. *Let $\xi = (E, p, B, F)$ be a G -stratified fiber bundle, H the isotropy of B . If F is an H -stratified pseudomanifold, then E is a G -stratified pseudomanifold.*

Proof. Fix a singular stratum S in E and a point $x \in S$. We must prove the existence of a link L_S depending only on S and, a distinguished slice (S_x, β, L_S) in x . For this purpose, let's take a trivializing chart

$$\varphi : U \times F \rightarrow p^{-1}(U) \in \mathcal{A}$$

such that $x \in p^{-1}(U)$. Take $z = p(x)$ and a G -slice V_z in B . Since V_z is contractible, we assume that $V_z \cong \mathbb{R}^k$ and V_z is contained in U .

Write $\varphi^{-1}(x) = (z, y) \in V_z \times F$ and take S' the stratum in F containing y . Since F is an H -stratified pseudomanifold, we can choose a distinguished slice S_y in y ; say

$$\beta_0 : S_y \rightarrow \mathbb{R}^i \times c(L_{S'})$$

Consider the following composition

$$\varphi(V_z \times S_y) \xrightarrow{\varphi^{-1}} V_z \times S_y \xrightarrow{\iota \times \beta_0} V_z \times \mathbb{R}^i \times c(L_{S'}) \cong \mathbb{R}^{i+k} \times c(L_{S'})$$

We will show that

$$(S_x, \beta, L_S) = (\varphi(V_z \times S_y), (\iota \times \beta_0) \circ \varphi^{-1}, L_{S'})$$

is a distinguished slice in x . We proceed in three steps.

• **L_S only depends on S :** If $(U', \psi) \in \mathcal{A}$ is another trivializing chart covering x , $\psi^{-1}(x) = (z, y') \in V_z \times F$ and $\beta'_0 : S_{y'} \rightarrow \mathbb{R}^i \times c(L_{S''})$ is a distinguished slice in y' ; then the composition $\beta'\beta^{-1}$ induces an H -isomorphism $L_S \xrightarrow{\cong} L_{S''}$.

• **S_x is a conical slice:** We verify the conditions (1) to (4) of §2.4.

- (1) Since V_z is a slice of $z \in B$, we have $gp(x) = p(gx) = p(x) \in V_z$ for any $g \in G_S$. So $G_S = H \cap G_S = H_S$, but φ is H -equivariant, hence $G_S = H_S = H_{S'}$. Again, since φ is H -equivariant and S_y is $H_{S'} = G_S$ invariant, we obtain that S_x is G_S -invariant.
- (2) Take $g \in G$, $x' \in S_x$ such that $gx' \in S_x$. Then $gp(x') = p(gx') \in V_z$, so $g \in H$ and $gp(x') = p(x')$. Since φ is H -equivariant, if $x' = \varphi(p(x'), y)$ then $g.x' = \varphi(p(x'), gy)$, and $gy \in S_y$; hence $g \in H_{S'} = G_S$.
- (3) Take a slice S_e of the identity element $e \in G$ with respect to the action of H . Since S_e is contractible, we can assume that $S_e V_z \subset U$. Notice that $S_e H$ is open in G . Since $GS_x = \bigcup_{g \in G} g(S_e H)S_x$, we only have to show that $(S_e H)S_x$ is open in X . But φ is H -equivariant and the action of H on V_z is trivial, so we get the following equality

$$(S_e H)S_x = S_e (H\varphi(V_z \times S_y)) = S_e \varphi(V_z \times HS_y)$$

Since HS_y is open in F we deduce that $S_e \varphi(V_z HS_y)$ is open in $S_e \varphi(V_z \times F)$. Finally we show that $S_e \varphi(V_z \times F) = S_e p^{-1}(V_z)$ is open in X : Since p is equivariant and $S_e V_z$ is open in U the set $S_e p^{-1}(V_z) = p^{-1}(S_e V_z) = p^{-1}(S_e H V_z)$ is open in $p^{-1}(U)$ (and so in X).

- (4) It is straightforward that the map β is a G_S -equivalence.

• **S_x is a distinguished slice:** We will show that usual the map

$$\alpha : G \times_{G_S} S_x \rightarrow X$$

is a (stratified) embedding.

- (a) **α preserves the strata:** Take a stratum S^0 in S_x . We will prove that $G'S^0$ is an open subset in some stratum of X , for any connected component $G' \subset G$. It is enough to prove this for the connected component G_0 of the identity element $e \in G$. Let H_0 be the connected component of the identity element $e \in H$. The set $S_e H_0$ is a connected open subset in $S_e H$, so is also connected and open in G_0 . Since $G_0 S^0$ is connected, we need to prove that $S_e H_0 S^0$ is open in some stratum of X . But $S_e H S_x$ is contained in $p^{-1}(S_e V_z)$ and φ is a stratified embedding, and so we only have to show that $\varphi^{-1}(S_e H_0 S^0)$ is open in some stratum of $(S_e V_z) \times F$. Consider the map

$$\begin{aligned} \Delta : S_e H \times V_z \times S_y &\rightarrow (S_e V_z) \times F \\ (gh, b, l) &\mapsto (ghb, (gh)_\varphi \varphi(b)(z)) = (gb, g_\varphi \varphi(b)(hz)) \end{aligned}$$

Let S^1 be the stratum of S_y such that $S^0 = \varphi(V_z \times S^1)$. By hypothesis S_y is a distinguished slice of y in F , and there is a stratum S^2 in F such that $H_0 S^1$ is open

in S^2 . Notice that

$$\varphi^{-1}(S_e H_0 S^0) = \Delta(S_e H_0 \times V_z \times S^1) = \Delta(S_e \times V_z \times H_0 S^1)$$

Also, since φ is H -equivariant, we have

$$p(\varphi^{-1}(S_e H_0 S^0)) = S_e V_z$$

Hence the projection $pr_2 : U \times F \rightarrow F$ sends $\varphi^{-1}(S_e H_0 S^0)$ on some open subset of S^2 . Notice that $\varphi^{-1}(S_e H_0 S^0)$ is connected, so

$$pr_2(\varphi^{-1}(S_e H_0 S^0)) = \bigcup_{(g,b) \in S_e \times V_z} g_{\varphi\varphi}(b)(H_0 S^1)$$

is a connected subset of F . Each $g_{\varphi\varphi}(b)$ is an H -equivariant stratified isomorphism; hence $g_{\varphi\varphi}(b)(H_0 S^1)$ is open in some stratum of F with the same dimension of S^2 . Since $e_{\varphi\varphi}(b)(H_0 S^1) = H_0 S^1 \subset S^2$, by §1.4-(2) the set $\bigcup_{(g,b) \in S_e \times V_z} g_{\varphi\varphi}(b)(H_0 S^1)$ is contained in S^2 .

(b) **α is smooth on each stratum:** Since $G \times_{G_x} S_x$ has the quotient stratification induced on $G \times S_x$ by the action of H , the stratification of S_x is induced by X and the action of G is smooth on each stratum of $G \times X$. We conclude that the restriction of α to each stratum is smooth. \square

3.3. Equivariant tubular neighborhoods An equivariant tubular neighborhood is a conical locally trivial fiber bundle. For a detailed introduction the reader can see [7], [11]. In [1], the tubular neighborhoods are used in order to show the existence of an unfolding for any manifold endowed with a Thom-Mather structure. We will provide an equivariant version of this fact for any G -stratified pseudomanifold.

Let X be a G -stratified pseudomanifold with $d(X) > 0$. Let's take a singular stratum S in X . An **equivariant tubular neighborhood** of S is a G -stratified fiber bundle $(T_S, \tau_S, S, c(L_S))$ with (maximal) trivializing atlas \mathcal{A} , verifying

- (1) T_S is an open invariant neighborhood of S and the inclusion $S \rightarrow T_S$ is a section of $\tau_S : T_S \rightarrow S$.
- (2) G preserves the conical radius: For any two charts $(U, \varphi), (U', \varphi') \in \mathcal{A}$ such that $U' \cap g^{-1}U \neq \emptyset$ for some $g \in G$, there is a map

$$g_{\varphi, \varphi'} : U' \cap g^{-1}U \rightarrow \text{Iso}_{G_S}(L_S, \mathcal{S}_{L_S})$$

such that

$$\varphi^{-1}g\varphi'(b, [l, r]) = (gb, [g_{\varphi, \varphi'}(b)l, r])$$

This allows us to define a (global) **radius** on T_S , as the map $\rho_S : T_S \rightarrow [0, \infty)$ satisfying

$$\rho_S(\varphi(z, [l, r])) = r \quad \forall (z, [l, r]) \in U \times c(L_S); (U, \varphi) \in \mathcal{A}$$

We also define the **radial action** $\delta_S : \mathbb{R}^+ \times T_S \rightarrow T_S$ as follows

$$\delta_S(r, x) = \varphi(z, [l, rt]) \quad \forall (z, [l, t]) \in U \times c(L_S); (U, \varphi) \in \mathcal{A} \quad (\text{for } x = \varphi(z, [l, t])).$$

We will write rx instead of $\delta_S(r, x)$ in the future. These functions satisfy

- (a) $\rho_S(rx) = r\rho_S(x)$ and $\rho_S(gx) = \rho_S(x)$ for any $r \in \mathbb{R}^+$, $x \in T_S$, $g \in G$.
- (b) $S \cap \rho_S^{-1}(0, \infty) = \emptyset$
- (c) The radial action commutes with the action of G on T_S .

3.4. Thom-Mather spaces (see [12], [13]): A **Thom-Mather G -stratified pseudomanifold** is a pair (X, \mathcal{T}) where X is a G -stratified pseudomanifold and $\mathcal{T} = \{T_S : S \in \mathcal{S}_X^{sing}\}$ is a family of equivariant tubular neighborhoods satisfying the following condition:

$$T_S \cap T_R \neq \emptyset \Leftrightarrow R \leq S \text{ or } S \leq R$$

for any two singular strata R, S in X . We will usually omit the family \mathcal{T} if there is no possible confusion.

3.5. Examples Here are some examples of G -stratified tubular neighborhoods.

- (1) Following [2, p.306], for any manifold M endowed with a smooth action $\Phi : G \times M \rightarrow M$ there is a riemannian metric μ such that G acts by μ -isometries. By the local properties of the exponential map, each singular stratum S of M has a smooth G -equivariant tubular neighborhood which can be realized as the normal fiber bundle $N_\mu(S)$ over S with respect to μ . The cocycles of this bundle are orthogonal actions. Hence, this tubular neighborhood is actually a G -stratified tubular neighborhood.
- (2) If L is a compact G -stratified pseudomanifold, the map $c(L) \rightarrow \{\star\}$ is a G -stratified tubular neighborhood of the vertex.
- (3) If $\xi = (T_S, \tau_S, S, c(L_S))$ is a G -stratified tubular neighborhood of S in X , then $(M \times T_S, \iota_M \times \tau_S, M \times S, c(L_S))$ is a G -stratified tubular neighborhood of $M \times S$ in $M \times X$; for any connected manifold M .
- (4) If $f : Y \rightarrow X$ is a G -equivariant isomorphism, then for any G -stratified tubular neighborhood $\xi = (T_S, S, \tau_S, c(L_S))$ of a stratum S in X ; the pull-back $f^*(\xi) = (f^{-1}(T_S), f^{-1}\tau_S f, f^{-1}(S), c(L_S))$ is a G -stratified tubular neighborhood of $f^{-1}(S)$ in Y .

Proposition 3.6. *Let X be a G -stratified pseudomanifold, K a closed subgroup of G . Write $\pi : X \rightarrow X/K$ for the orbit map induced by the action of K . Let $\xi = (T_S, \tau_S, S, c(L_S))$ be an equivariant tubular neighborhood of S in X and write*

$$\overline{\tau_S} : \pi(T_S) \rightarrow \pi(S)$$

for the induced quotient map. Then $\xi/K = (\pi(T_S), \overline{\tau_S}, \pi(S), c(L_S/G_S \cap K))$ is an equivariant tubular neighborhood of $\pi(S)$ in X/K .

Proof. Since π is an open map, $\pi(T_S)$ is an open neighborhood of $\pi(S)$ in X/K . Also the inclusion $\pi(S) \rightarrow \pi(T_S)$ is a section of $\overline{\tau_S} : \pi(T_S) \rightarrow \pi(S)$. In order to prove that ξ/K is a G -stratified tubular neighborhood we should first verify that it is a G -stratified fiber bundle, but the conditions §3.1-(1) to (4) are straightforward.

Now we will prove §3.3-(2), which implies §3.1-(5). We will show that the trivializing atlas $\mathcal{A} = \{(U, \varphi)\}$ of ξ induces a trivializing atlas $\mathcal{A}/K = \{(V, \psi)\}$ of ξ/K .

Write $\pi' : L_S \rightarrow L_S/G_S \cap K$ for the orbit map induced by the action of $G_S \cap K$ in L_S .

• **Trivializing charts:** Take a chart $(U, \varphi) \in \mathcal{A}$ and a point $x \in U$. Take also a K -slice V of x in S , we assume that $V \subset U$. Since G_S acts trivially on V and KV is open in S we deduce that

$$V = V/G_S \cap K = \pi(KV)$$

is open in $\pi(S)$. Since φ is G_S -equivariant, the function

$$(1) \quad \psi : V \times c(L_S/G_S \cap K) \rightarrow \pi(T_S) \quad \psi(b, [\pi'(l), r]) = \pi(\varphi(b, [l, r]))$$

is well defined. Moreover, ψ is injective because G acts by isomorphisms and V is a K -slice in S . Notice that $W = KV \cap U$ is open in U ; since G also preserves the radius in T_S ,

$$\text{Im}(\psi) = \pi(\varphi(W \times c(L_S)))$$

Hence $\text{Im}(\psi)$ is open in X/K . It is straightforward that ψ sends smoothly strata onto strata, so actually ψ is an embedding.

• **Atlas and cocycles:** We consider the family $\mathcal{A}/K = \{V, \psi\}$ of all the pairs (V, ψ) as in (1). We will show that \mathcal{A}/K is a trivializing atlas of ξ/K . Take two charts $(V, \psi); (V', \psi') \in \mathcal{A}/K$ respectively induced by $(U, \varphi); (U', \varphi') \in \mathcal{A}$. Assume that there is some $\overline{g_0} \in G/K$ such that $\overline{g_0}^{-1}V \cap V' \neq \emptyset$; so $g^{-1}U \cap U' \neq \emptyset$ for some $g \in g_0K$. By §3.3-(2), there is a map

$$g_{\varphi\varphi'} : g^{-1}U \cap U' \rightarrow \text{Iso}_{G_S}(L_S, \mathcal{S}_{L_S})$$

satisfying

$$g\varphi'(b, [l, r]) = \varphi(gb, [g_{\varphi\varphi'}(b)(l), r]) \quad (b, [l, r]) \in (g^{-1}U \cap U') \times c(L_S)$$

Passing to the orbit space $L_S/G_S \cap K$ we obtain the induced map

$$\overline{g_0}\psi\psi' : \overline{g_0}^{-1}V \cap V' \rightarrow \text{Iso}_{(G_S/G_S \cap K)}(L_S/G_S \cap K, \mathcal{S}_{L_S/G_S \cap K})$$

satisfying

$$\overline{g_0}\psi'(b, [\pi'(l), r]) = \psi(\overline{g_0}b, [\overline{g_0}\psi\psi'(b)(\pi'(l)), r]); \quad (b, [\pi'(l), r]) \in (\overline{g_0}^{-1}V \cap V') \times c(L_S/G_S \cap K)$$

Notice that, by definition, G/K preserves the radius of $\pi(T_S)$. \square

4. EQUIVARIANT UNFOLDINGS

An unfolding of a stratified pseudomanifold is an auxiliary construction which allows us to define the intersection cohomology from the point of view of differential forms [1],[3]. For a detailed introduction to unfoldings, the reader can see [4], [9]. In this section we introduce equivariant unfoldings, these are a suitable adaptation of the usual unfoldings to the equivariant category. We also show that for any G -manifold, considered as a G -stratified pseudomanifold, there is always an equivariant unfolding which induces a canonical unfolding on the orbit space.

4.1. Equivariant unfoldings Broadly speaking, an un unfolding of a stratified pseudomanifold X is a manifold \tilde{X} and a surjective continuous map $\mathbb{L} : \tilde{X} \rightarrow X$ such that $\mathbb{L}^{-1}(X - \Sigma)$ is a union of finitely many disjoint copies of $X - \Sigma$, and which smoothly unfolds the singular part so that the restriction $\mathbb{L} : \mathbb{L}^{-1}(S) \rightarrow S$ is a submersion, for any singular stratum S .

As for the usual unfoldings, the definition of an equivariant unfolding is made by induction on the depth. Let X be a G -stratified pseudomanifold. An **equivariant unfolding** of X is a manifold \tilde{X} together with a smooth free action $\tilde{\Phi} : G \times \tilde{X} \rightarrow \tilde{X}$; a surjective, continuous, equivariant map

$$\mathbb{L} : \tilde{X} \rightarrow X$$

and a family of equivariant unfoldings $\{\mathbb{L}_S : \tilde{L}_S \rightarrow L_S\}_{S \in S_X^{sing}}$ where S runs on the singular strata of X ; satisfying:

- (1) The restriction $\mathbb{L} : \mathbb{L}^{-1}(X - \Sigma) \rightarrow X - \Sigma$ is a smooth finite trivial covering.
- (2) For each singular stratum S and each $x \in S$, there is a **liftable modelled chart**, i.e.; a commutative square

$$\begin{array}{ccc} U \times \tilde{L}_S \times \mathbb{R} & \xrightarrow{\tilde{\varphi}} & \tilde{X} \\ \mathbb{L}_c \downarrow & & \downarrow \mathbb{L} \\ U \times c(L_S) & \xrightarrow{\varphi} & X \end{array}$$

such that

- (a) (U, φ) is a G_S -equivariant chart of x modelled on L_S .
- (b) $\tilde{\varphi}$ is a G_S -equivariant smooth embedding on an open subset of \tilde{X} .
- (c) The map \mathbb{L}_c is given by the rule $\mathbb{L}_c(u, z, t) = (u, [\mathbb{L}_{L_S}(z), |t|])$.

A G -stratified pseudomanifold X is said to be **unfoldable** whenever it has an equivariant unfolding.

4.2. Examples Here are some examples of equivariant unfoldings.

- (1) For any free smooth action $\Phi : G \times M \rightarrow M$ the identity $\iota : M \rightarrow M$ is an equivariant unfolding.
- (2) If $\mathbb{L} : \tilde{X} \rightarrow X$ is an equivariant unfolding, then for any manifold M the product $\iota : M \times \tilde{X} \rightarrow M \times X$ is also an equivariant unfolding.
- (3) For any equivariant unfolding $\mathbb{L} : \tilde{L} \rightarrow L$ over a compact G -stratified pseudomanifold L , the map $\mathbb{L}_c : \tilde{L} \times \mathbb{R} \rightarrow c(L)$ defined above is also an equivariant unfolding.

4.3. Elementary unfolding of a G -stratified pseudomanifold The elementary unfolding of a Thom-Mather space is essentially the resolution of singularities given in [4] for the smooth case. This topological operation can be done because the stratification is controlled through a family of tubular neighborhoods. Under certain conditions, after the iterated composition of finitely many elementary unfoldings,

one obtains an equivariant unfolding as defined above. We follow the exposition of [1].

Henceforth we fix a Thom-Mather G -stratified pseudomanifold X , a closed (hence minimal) stratum S in X and an equivariant tubular neighborhood $(T_S, \tau_S, S, c(L_S))$ of S . Define the **unitary sub-bundle** as the set $E_S = \rho_S^{-1}(1)$; this is by construction a G -invariant stratified subspace of X . The restriction $\tau_S : E_S \rightarrow S$ is a G -stratified fiber bundle with fiber L_S . Consider the map

$$(2) \quad \mathbf{L}_{T_S} : E_S \times \mathbb{R} \rightarrow T_S \quad \mathbf{L}_{T_S}(x, t) = \begin{cases} |t| * x & \text{si } t \neq 0 \\ \tau_S(x) & \text{si } t = 0 \end{cases}$$

Each chart (U, φ) in the trivializing atlas provides a local description of \mathbf{L}_{T_S} through the following commutative square

$$\begin{array}{ccc} U \times L_S \times \mathbb{R} & \xrightarrow{\widehat{\varphi}} & E_S \times \mathbb{R} \\ \downarrow 1_U \times \mathbf{L}_C & & \downarrow \mathbf{L}_{T_S} \\ U \times cL_S & \xrightarrow{\varphi} & T_S \end{array}$$

where $\widehat{\varphi}(x, l, t) = (\varphi(x, [l, 1], t))$ and $\mathbf{L}_C(l, t) = [l, |t|]$. We also obtain the following properties:

- (a) The map $\widehat{\varphi}$ is a G_S -equivariant embedding.
- (b) The composition $\tau_S \circ \mathbf{L}_{T_S} : E_S \times \mathbb{R} \rightarrow S$ is a locally trivial fiber bundle with fiber $L_S \times \mathbb{R}$ and structure group $\text{Iso}_{G_S}(L_S, \mathcal{S}_{L_S})$.
- (c) $d(E_S \times \mathbb{R}) = d(E_S) = d(T_S) - 1$.

Now take a disjoint family of equivariant tubular neighborhoods $\{T_S : S \in \mathcal{S}_X^{\min}\}$ of the minimal strata. The **elementary unfolding** of X with respect to the family $\{T_S : S \in \mathcal{S}_X^{\min}\}$ is the pair $(\widehat{X}, \mathbf{L})$ constructed as follows: First \widehat{X} is the amalgamated sum

$$(3) \quad \widehat{X} = \left[\bigsqcup_{S \in \mathcal{S}_X^{\min}} E_S \times \mathbb{R} \right] \bigcup_{\theta} [(X - \Sigma^{\min}) \times \{\pm 1\}]$$

where θ is the map

$$(4) \quad \theta : \bigsqcup_{S \in \mathcal{S}_X^{\min}} E_S \times \mathbb{R}^* \rightarrow [X - \Sigma^{\min}] \times \{\pm 1\} \quad \theta(x, t) = (|t| * x, |t|^{-1}t)$$

Second, \mathbf{L} is the continuous map given by the rule

$$(5) \quad \mathbf{L} : \widehat{X} \rightarrow X \quad \mathbf{L}(x) = \begin{cases} \mathbf{L}_{T_S}(x) & x \in E_S \times \mathbb{R} \\ y & x = (y, j) \in (X - \Sigma^{\min}) \times \{\pm 1\} \end{cases}$$

Here there are some properties of the elementary unfoldings.

Proposition 4.4. *Let $\mathbf{L} : \widehat{X} \rightarrow X$ be the elementary unfolding of a Thom-Mather G -stratified pseudomanifold X . Then*

- (1) \widehat{X} is a G -stratified pseudomanifold, whose stratification is the family $\mathcal{S}_{\widehat{X}}$ consisting of all the following sets

$$\widehat{R} = \bigsqcup_{S \in \mathcal{S}_X^{\min}} (E_S \cap R) \times \mathbb{R} \bigsqcup_{\theta} (R \times \{\pm 1\})$$

where R runs over the non closed strata in X . Moreover, \widehat{X} satisfies the Thom-Mather condition.

- (2) The map L is a G -equivariant morphism. The restriction

$$L : L^{-1}(X - \Sigma^{\min}) \rightarrow X - \Sigma^{\min}$$

is a (trivial) double covering.

- (3) $d(\widehat{X}) = d(X) - 1$. In particular, if $d(X) = 1$ then $L : \widehat{X} \rightarrow X$ is an equivariant unfolding.
- (4) If X is compact, then so is \widehat{X} .
- (5) For any closed subgroup $K \subset G$, the induced map $\overline{L} : \widehat{X}/K \rightarrow X/K$ is an elementary unfolding.

Proof. (1) The stratification of \widehat{X} can be seen in [1]. Since each equivariant tubular neighborhood is a G -stratified pseudomanifold (because they are invariant open subsets of X); so are the unitary sub-bundles (see §3.2), and hence \widehat{X} is a G -stratified pseudomanifold. Now we verify the Thom-Mather condition: Take a family $\{T_S : S \in \mathcal{S}_X\}$ of equivariant tubular neighborhoods in X . Take also a stratum \widehat{R} in \widehat{X} induced by a non closed stratum R in X . Define

$$T_{\widehat{R}} = \bigsqcup_{S \in \mathcal{S}_X^{\min}} (E_S \cap T_R) \times \mathbb{R} \cup (T_R \times \{\pm 1\}) = L^{-1}(T_R)$$

where θ is the map given in the equation (4) of §4.3. This $T_{\widehat{R}}$ is an equivariant tubular neighborhood of \widehat{R} in \widehat{X} ; we leave the details to the reader.

(2) and (3) are straightforward, see again [1] for more details. The last observation of (3) is a consequence of def. §4.1.

(4) Since X is compact, \mathcal{S}_X^{\min} is finite. But \widehat{X} is the quotient of the finite family of compact spaces $\bigsqcup_{S \in \mathcal{S}_X^{\min}} (E_S \times [-1, 1])$ and $[X - \bigsqcup_{S \in \mathcal{S}_X^{\min}} \rho_S^{-1}[0, 1/2)] \times \{-1, 1\}$. Then

we get the result.

- (5) This is a consequence of §3.6. □

♣ *Remark 4.5.* With tubular neighborhood of 3.5-3, $\widehat{M \times X} = M \times \widehat{X}$, for any manifold M .

5. ITERATION OF ELEMENTARY UNFOLDINGS

From now on, we fix a Thom-Mather G -stratified pseudomanifold X . We will study the composition of finitely many elementary unfoldings, starting at X . As we have already seen, for any elementary unfolding $L : \widehat{X} \rightarrow X$, the space \widehat{X} is again

a Thom-Mather G -stratified pseudomanifold and satisfies $d(\widehat{X}) = d(X) - 1$. This allows us to ask for the behavior of a chain

$$(6) \quad X_l \xrightarrow{\mathbb{L}_l} X_{l-1} \xrightarrow{\mathbb{L}_{l-1}} \dots \xrightarrow{\mathbb{L}_2} X_1 \xrightarrow{\mathbb{L}_1} X$$

of elementary unfoldings, where $l = d(X)$. As we shall see, under certain conditions on the tubular neighborhoods, this iterative process leads us to an equivariant unfolding

$$\mathbb{L} : \widetilde{X} \rightarrow X$$

where $\widetilde{X} = X_l$ and $\mathbb{L} = \mathbb{L}_1 \dots \mathbb{L}_l$.

Recall the definition of a **saturated subspace** [1]. Let $Y \subset X$ be a stratified subspace of X . We say that Y is **saturated** whenever

$$Y \cap T_S = \tau_S^{-1}(Y \cap S) \quad \forall S \in \mathcal{S}_X$$

For instance, if S is a singular stratum and $U \subset S$ is open, then $Y = \tau_S^{-1}(U)$ is a saturated. Also the unitary sub-bundle $Y = E_S$ is saturated.

5.1. Transverse morphisms Now we introduce the family of transverse morphisms, whose main feature is the preservation of the tubular neighborhoods. Let $H \subset G$ be a closed subgroup, Y a Thom-Mather H -stratified pseudomanifold and M be a connected manifold. A morphism

$$\psi : M \times Y \rightarrow X$$

is **transverse** whenever:

- (1) $\text{Im}(\psi)$ is a saturated open subspace of X .
 - (2) If $\psi(M \times S) \subset R$ then $\psi^{-1}(T_R) = M \times T_S$, for any $R \in \mathcal{S}_X, S \in \mathcal{S}_Y$.
- Now let $\psi : M \times Y \rightarrow X$ be a transverse morphism. The **lifting** of ψ is, by definition, the map

$$\widehat{\psi} : M \times \widehat{Y} \rightarrow \widehat{X} \quad \widehat{\psi}(m, z, t) = \begin{cases} (\psi(m, z), t) & (m, z, t) \in M \times E_S \times \mathbb{R} \\ (\psi(m, z), t) & (m, z, t) \in M \times (Y - \Sigma^{\min}) \times \{\pm 1\} \end{cases}$$

This is the unique morphism such that the diagram

$$\begin{array}{ccc} M \times \widehat{Y} & \xrightarrow{\widehat{\psi}} & \widehat{X} \\ \downarrow \iota_M \times \mathbb{L}_Y & & \downarrow \mathbb{L}_X \\ M \times Y & \xrightarrow{\psi} & X \end{array}$$

commutes.

5.2. Examples For any smooth effective action of G in a manifold, the trivializing charts of the tubular neighborhoods are transverse morphisms. In order to see this, take a manifold M endowed with a smooth action $\Phi : G \times M \rightarrow M$ and an invariant metric μ in M . Recall that M has a natural structure of Thom-Mather G -stratified pseudomanifold, where \mathcal{S}_M is the stratification induced by the orbit types of the

action. For any singular stratum S with codimension $\text{codim}(S) = q + 1 > 0$, the equivariant tubular neighborhood $T_S = N_\mu(S)$ is the normal fiber bundle over S induced by μ (see §3.5). Take also a trivializing chart

$$\varphi : U \times c(\mathbb{S}^q) \rightarrow \tau_S^{-1}(U)$$

We claim that φ is transverse. First notice that $\text{Im}(\varphi)$ is a saturated open subspace in M , so we only have to verify §5.1-(2). Let S' be a stratum in $c(\mathbb{S}^q)$, R a stratum in M . Suppose that $\varphi(U \times S') \subset R$. We consider the following cases:

- $S' = \{\star\}$ is the vertex: It is straightforward, since $R = S$ and $T_{S'} = c(\mathbb{S}^q)$.
- $S' = S'' \times \mathbb{R}^+$ for some stratum S'' in \mathbb{S}^q : Then $S < R$. We consider in T_S the following decomposition of the metric:

$$\mu|_{T_S} = \mu_H + \mu_V$$

corresponding to the orthogonal decomposition of the tangent $T(T_S)$ in the horizontal and vertical subfiber bundles. Hence

$$\varphi^{-1}(T_R) = \varphi^*(N_\mu(R)) = N_{\varphi^*(\mu)}(U \times S') = U \times N_{\mu_V}(S') = U \times T_{S'}$$

Now we show two easy properties of the transverse morphisms.

Proposition 5.3. *Let K, H a closed subgroups of G , L a Thom-Mather H -stratified pseudomanifold, $\psi : M \times L \rightarrow X$ a transverse morphism. Then*

- (1) *The lifting $\hat{\psi} : M \times \hat{L} \rightarrow \hat{X}$ is transverse.*
- (2) *The induced quotient map $\bar{\psi} : M \times (L/H \cap K) \rightarrow X/K$ is transverse.*

Proof. (1) is straightforward from the def. §5.1. (2) is a consequence of §3.6. \square

Finally, we provide a sufficient condition for the existence of an equivariant unfolding, depending on the transversality of the tubular neighborhoods.

Theorem 5.4. *Let X be a Thom-Mather G -stratified pseudomanifold. Suppose that for any singular stratum S , each trivializing chart*

$$\varphi : U \times c(L_S) \rightarrow T_S$$

is transverse. Then

- (1) *The composition of the l elementary unfoldings of starting at X induces an equivariant unfolding $L : \tilde{X} \rightarrow X$ where \tilde{X} is the last (non trivial) elementary unfolding and $L = L_1 L_2 \dots L_l$ (see eq. (6) at the beginning of this section).*
- (2) *For any closed subgroup $K \subset G$, the induced map $\bar{L} : \tilde{X}/K \rightarrow X/K$ is an unfolding.*

Proof. (1) Take a family of equivariant tubular neighborhoods in X with transverse trivializing charts. Let

$$X_l \xrightarrow{L_l} X_{l-1} \xrightarrow{L_{l-1}} \dots \xrightarrow{L_2} X_1 \xrightarrow{L_1} X$$

be the chain of elementary unfoldings induced by this family of tubular neighborhoods. We proceed by induction on $l = d(X)$; for $l = 1$ the statements are trivial.

Suppose that $l > 1$ and assume the inductive hypothesis, so $L' : \tilde{X} \rightarrow X_1$ is an equivariant unfolding, for $\tilde{X} = X_l$ and $L' = L_2 \dots L_l$. Take a closed stratum S and a transverse trivializing chart

$$\varphi : U \times c(L_S) \rightarrow \tau_S^{-1}(U) \subset T_S$$

Apply the chain of elementary unfoldings and use §5.3; you will get the following commutative diagram:

$$\begin{array}{ccc} U \times \widetilde{L}_S \times \mathbb{R} & \xrightarrow{\widetilde{\psi}_1} & \tilde{X} \\ \downarrow \iota_U \times L_S \times \iota_{\mathbb{R}} & & \downarrow L' \\ U \times L_S \times \mathbb{R} & \xrightarrow{\psi_1} & X_1 \\ \downarrow \iota_U \times L_S & & \downarrow L_1 \\ U \times c(L_S) & \xrightarrow{\psi} & X \end{array}$$

We conclude that $L = L_1 L' : \tilde{X} \rightarrow X$ is an equivariant unfolding.

(2) This is a consequence of §4.4-(6). □

Corollary 5.5 (Unfolding of a G -manifold).

Let M be a manifold, $\Phi : G \times M \rightarrow M$ a smooth effective action, possibly with fixed points. Endow M with the stratification induced by the orbit types and the usual structure of a Thom-Mather G -stratified pseudomanifold. Then there is an equivariant unfolding $L : \tilde{M} \rightarrow M$.

Proof. Apply the above theorem to the transverse charts obtained in §5.2. □

ACKNOWLEDGMENTS

I would like to thank M. Saralegi for some helpful conversations, and R. Popper for some accurate remarks. While writing this work, I received the hospitality of the staff in the Math Department, Université D'Artois; and the financial support of the ECOS-Nord project and the CDCH-Universidad Central de Venezuela.

REFERENCES

- [1] BRASSELET, J.P., G. HECTOR et M. SARALEGI, *Théorème de De Rham Pour les variétés stratifiées* Ann. Global Anal. Geom. #9, 201-243 (1991).
- [2] BREDON, G. *Introduction to Compact Transformation Groups*. Pure and Appl. Math. vol. #46 (1972) Academic Press, New York and London.
- [3] BRYLINSKY, J. L. *Equivariant Intersection Cohomology*. Contemporary Math. **132** (1992), 5-32.
- [4] DAVIS, M. *Smooth G -Manifolds as collections of Fiber Bundles*. Pacific J. Math. #77 (1978), 315-363.
- [5] HAEFLIGER, A. "Introduction to piecewise linear intersection homology" (Bern, 1983)" in *Intersection Cohomology-Swiss seminars*. (Bern, 1983). Progress in Mathematics Vol.50, 1-22. Birkhäuser. Boston (1984).

- [6] PADILLA, G. *Intersection cohomology of S^1 -actions on pseudomanifolds*. (2003) Submitted preprint.
- [7] PFLAUM, M. *Analytic and Geometric study of Stratified Spaces*. (2000) Springer-Verlag.
- [8] POPPER, R. *Compact Lie Groups Acting on Pseudomanifolds*. Illinois J. Math., #**44** (2000), 1-19.
- [9] SARALEGI, M. *Homological Properties of Stratified Spaces*. Illinois J. Math. #**38** (1994), 47-70.
- [10] STEENROD, N. *The topology of fiber bundles*. Princeton University Press. Princeton-New Jersey . (1951).
- [11] THOM, R. *Ensembles et morphismes stratifiés*. Bull. Amer. Math. Soc. **75**, 240-284 (1969).
- [12] VERONA, A. *Le théoreme de DeRham pour les préstratifications abstraites*. C.R. Acad. Sci. Paris **273** (1971), 886-889.
- [13] VERONA, A. *Stratified mappings, structure and triangulability*. Lecture Notes in Math. Springer-Verlag. Vol. **1102**. Springer-Verlag. New York-Heidelberg- Berlin (1984).

ESCUELA DE MATEMATICAS, FACULTAD DE CIENCIAS, UCV, CARACAS VENEZUELA
E-mail address: dalmagro@euler.ciens.ucv.ve